PTSD CheckList – Civilian Version (PCL-C)

Below is a list of problems and complaints that people sometimes have in response to stressful life experiences. Please read each one carefully, pick the answer that indicates how much you have been bothered by that problem *in the last month*.

No.	Response	Not at all (1)	A little bit (2)	Moderately (3)	Quite a bit (4)	Extremely (5)
1.	Repeated, disturbing <i>memories</i> , <i>thoughts</i> , <i>or images</i> of a stressful experience from the past?					
2.	Repeated, disturbing <i>dreams</i> of a stressful experience from the past?					
3.	Suddenly <i>acting</i> or <i>feeling</i> as if a stressful experience <i>were happening</i> again (as if you were reliving it)?					
4.	Feeling <i>very upset</i> when <i>something</i> <i>reminded</i> you of a stressful experience from the past?					
5.	Having <i>physical reactions</i> (e.g., heart pounding, trouble breathing, or sweating) when <i>something</i> <i>reminded</i> you of a stressful experience from the past?					
6.	Avoid <i>thinking about</i> or <i>talking</i> <i>about</i> a stressful experience from the past or avoid <i>having feelings</i> related to it?					
7.	Avoid <i>activities</i> or <i>situations</i> because they <i>remind you</i> of a stressful experience from the past?					
8.	Trouble <i>remembering important</i> <i>parts</i> of a stressful experience from the past?					
9.	Loss of <i>interest in things that you used to enjoy?</i>					
10.	Feeling <i>distant</i> or <i>cut</i> off from other people?					
11.	Feeling <i>emotionally numb</i> or being unable to have loving feelings for those close to you?					
12.	Feeling as if your <i>future</i> will somehow be <i>cut short</i> ?					
13.	Trouble <i>falling</i> or <i>staying asleep</i> ?					
14.	Feeling <i>irritable</i> or having <i>angry outbursts</i> ?					
15.	Having difficulty concentrating?					
16.	Being " <i>super alert</i> " or watchful on guard?					
17.	Feeling <i>jumpy</i> or easily startled?					

PCL-M for DSM-IV (11/1/94) Weathers, Litz, Huska, & Keane National Center for PTSD-Behavioral Science Div.

How is the PCL Scored?

1) Add up all items from each of the 17 items for a total severity score (range = 17-85) 17-29 This cut off shows little to no severity.

28-29 Some PTSD symptoms - If you are seeing or will be seeing a therapist, print the results of this Quiz and take to your therapist for further evaluation.

30–44 Moderate to Moderately High severity of PTSD symptoms - If you are seeing or will be seeing a therapist, print the results of this Quiz and take to your therapist for further evaluation.

45-85 High Severity of PTSD symptoms - If you are seeing or will be seeing a therapist, print the results of this Quiz and take to your therapist for further evaluation.